

LA-UR-12-25958

Approved for public release; distribution is unlimited.

Title: Berezinskii-Kosterlitz-Thouless Transition in Quantum Critical Metals

Author(s): She, Jianhuang

Intended for: talk in Munich and Leiden



Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.


Berezinskii-Kosterlitz-Thouless Transition in Quantum Critical Metals

Jian-Huang She


Los Alamos National Laboratory

JHS & A. V. Balatsky, Phys. Rev. Lett. 109,077002(2012)


Lower dimension, stronger correlation, higher T_c ?




MgB_2 : $T_c = 39\text{K}$





$T_c^{\text{max}} = 93\text{K}$



$\text{Ba}_{0.6}\text{K}_{0.4}\text{Fe}_2\text{As}_2$: 38K



$\text{SmFeAsO}_{0.85}$: 55K



CePt_2In_7
 12 of 14 members of
 $\text{Ce}_m\text{M}_n\text{In}_{3m+2n}$
 and PuMGA_5 family
 are superconductors

Increasing T_c
 100x

Eric Bauer

Lower dimension, more fluctuation, lower T_c ?

- 1, High dimension: mean field behavior ($d > \text{upper critical dimension}$)
- 2, Low dimension, more fluctuations (Mermin-Wagner theorem) :
continuous global symmetry can not be broken in 2d at finite T.


$$H \sim \int d^d x (\nabla \phi)^2 \qquad \langle \phi^2(0) \rangle \sim \int \frac{d^d k}{(2\pi)^d} \frac{1}{k^2}$$

$d=1,2$, the integral diverges at IR. ($d=2$: lower critical dimension)


How to get high T_c (nature's solution): **heterostructure**

2 dimensional correlation, 3 dimensional phase transition


Strongly correlated heterostructures: oxide interface



Haraldsen, Wolfle, Balatsky, PRB 2012



Cavaglia *et al.*, Nature 2008





Hwang *et al.* Nat. Mat. 2012


How about f-electrons?

Heavy fermion goes two dimensional: CeIn₃/LaIn₃


Molecular Beam Epitaxy



Shishido *et al.* Science 2010




Dimensionally tuned quantum criticality





Heavy fermion goes two dimensional: CeCoIn₅/YbCoIn₅

Heavy-fermion superconductor
CeCoIn₅ ($T_c=2.3\text{K}$)




Molecular Beam Epitaxy
(MBE)





Mizukami *et al.* Nat. Phys. 2011

Two dimensional Kondo lattice





Mizukami *et al.* Nat. Phys. 2011





Thompson, Nat. Phys. 2011




Superconductivity in HF superlattices



With magnetic field



2d or 3d: proximity effect




Thickness of leakage region much larger than interlayer spacing

$$l = \frac{\hbar v_N}{2\pi k_B T} \simeq 100\text{nm}$$

$$d \simeq 3.7\text{nm}$$

Suppressed proximity effect from mass mismatch

E. W. Fenton (1985)



Transmitted probability current

$$\lambda \approx \frac{4m_{\text{light}}}{m_{\text{heavy}}} \quad m_h \approx 10^2 m_l$$

Transmission: one percent !

Valls, Bryan, Zutic (2010)




FIG. 3: (Color online) The spatial dependence of the pair amplitude $F(Z)$ for five different mismatch values $\Lambda = 1/4$ (purple), $1/2$ (green), 4 (cyan), 2 (blue), and 1 (red), from top to bottom on the right side. The results are given for $H_B = 0$, $g = -1/3$ at $T = 0.3T_c$.

$$\text{Effective barrier: } Z = (Z_0^2 + (1 - r)^2 / 4r)^{1/2}$$

$$r = v_S / v_N$$

Blonder, Tinkham, Klapwijk (1982)


Two dimensional superconductor



N




N





N



N



Gap and Tc



Pauli limited upper critical field

$$H_{c2}^{\text{Pauli}} = \sqrt{2} \Delta / g \mu_B$$

Phase fluctuations!

This is a Berezinskii-Kosterlitz-Thouless transition!



Consider a system with $U(1)$ symmetry, e.g. XY-magnet, superfluid...

Landau: symmetry breaking, second-order phase transition

$$\text{As } |\mathbf{r}_1 - \mathbf{r}_2| \rightarrow \infty, \langle \Phi(\mathbf{r}_1) \Phi(\mathbf{r}_2) \rangle \sim \begin{cases} e^{-|\mathbf{r}_1 - \mathbf{r}_2|/\xi}, & \text{for } T > T_c \\ \text{constant}, & \text{for } T < T_c \end{cases}$$

Mermin-Wagner: no broken continuous symmetry for $d=2,1$, due to strong fluctuations

BKT: quasi-long-range-order and true phase transition in 2d

$$\text{As } |\mathbf{r}_1 - \mathbf{r}_2| \rightarrow \infty, \langle \Phi(\mathbf{r}_1) \Phi(\mathbf{r}_2) \rangle \sim \left(\frac{|\mathbf{r}_1 - \mathbf{r}_2|}{r_0} \right)^\eta, \text{ for } T < T_c.$$

Vortex: topological defect in systems with U(1) symmetry


Superconducting order parameter: $\Psi(\mathbf{r}, t) \equiv A(\mathbf{r}, t) \exp i\phi(\mathbf{r}, t)$

Decompose phase into smooth and singular parts:

$$\phi(\mathbf{r}) = \phi_{\text{sm}}(\mathbf{r}) + \phi_{\text{sg}}(\mathbf{r})$$

$$\oint d\mathbf{r} \cdot \nabla \phi_{\text{sm}}(\mathbf{r}) = 0$$

$$\oint d\mathbf{r} \cdot \nabla \phi_{\text{sg}}(\mathbf{r}) = 2\pi n$$




Vortex-vortex interaction



$$\frac{\mathcal{H}_v}{k_B T} = -\pi K \sum_{ij} n_i n_j \log \frac{|\mathbf{r}_i - \mathbf{r}_j|}{R_0} - \log y \sum_i n_i^2$$


Stiffness: $K = n_s \hbar^2 / 4mk_B T$

Fugacity: $y = e^{-E_c/k_B T}$



High T

free vortex



Low T

bound pairs

UNCLASSIFIED

Slide 16

Feature I: universal jump in superfluid density

Kosterlitz's RG equation:


$$\frac{d}{dl} K^{-1}(l) = 4\pi^3 y^2(l),$$

$$\frac{d}{dl} y(l) = [2 - \pi K(l)] y(l)$$

Superfluid density has a jump at T_c :


$$K \equiv \frac{\hbar^2 \rho_s(T)}{m^2 k_B T} = \begin{cases} 0, & \text{for } T = T_c^+ \\ 2/\pi, & \text{for } T = T_c^- \end{cases}$$

Mean field transition: $\rho_s(T) \sim 1 - \left(\frac{T}{T_c}\right)^\alpha$



Bonn, 2006

Superfluid helium film: $\Delta P \propto \rho_s$



McQueeney, Agnolet, Reppy (1984)

Feature II: thickness dependence of transition temperature

Universal relation: $k_B T_{\text{BKT}} = \frac{\pi \hbar^2 n_s^{2D}(T_{\text{BKT}})}{8m\epsilon_c}$


Carrier density: $n_s^{2D} = n_s^{3D} d$

$$n_s^{3D}(T) = n_s^{3D}(0) \lambda_b^2(0) / \lambda_b^2(T)$$

Penetration depth: $\lambda(T) = \frac{\lambda(0)}{\sqrt{1 - (T/T_{c0})^\alpha}}$

Thickness dependence of BKT temperature:

$$\frac{T_{\text{BKT}}}{1 - (T_{\text{BKT}}/T_{c0})^\alpha} = \frac{\pi \hbar^2 n_s^{3D}(0)}{8k_B m \epsilon_c} d$$




Matsuda *et al.*, 1993

Phase fluctuations in cuprates

Uemura relation (underdoped): $T_c = \text{const} \times \rho_s(0)$

$$H = \frac{1}{2} m^* n_s(0) \int dr v_s^2 \quad v_s = \hbar \nabla \theta / 2m^*$$

$$T_\theta^{\text{max}} = A V_0 \quad V_0 = \frac{(\hbar c)^2 a}{16\pi e^2 \lambda^2(0)}$$



Emery, Kivelson, 1995

Homes' law for cuprates (all doping): $\rho_s \propto \sigma_{dc} T_c$

Towards a holographic realization of Homes' law

Johanna Erdmenger, Patrick Kerner and Steffen Müller

*Max-Planck-Institute for Physics, Werner Heisenberg Institut,
80805 Munich, Germany*

Feature III: temperature dependence of resistivity

Above BKT temperature, there are unpaired vortices, which give rise to resistivity.

Kosterlitz's RG equation:



$$\begin{aligned}\frac{d}{dl}K^{-1}(l) &= 4\pi^3 y^2(l), \\ \frac{d}{dl}y(l) &= [2 - \pi K(l)]y(l)\end{aligned}$$

Correlation length: $\xi(T > T_c) \sim e^{b/\sqrt{T-T_c}}$






Density of unpaired vortices: $n_f \propto \xi^{-2}$

Resistivity from SC fluctuations: $\rho_{fl} \propto n_f \propto e^{-2b/\sqrt{T-T_c}}$

Check: are they BKT transitions?





Resistivity (feature III)



$$\rho(T) = \rho_0 e^{-b(T-T_{KT})^{-1/2}}$$


$$\left[\frac{d \ln \rho(T)}{dT} \right]^{-2/3} = \left(\frac{2}{b} \right)^{2/3} (T - T_{KT})$$

BKT transition temperature (feature II)



$$\frac{T_{\text{BKT}}}{1 - (T_{\text{BKT}}/T_{c0})^2} = \frac{\pi \hbar^2 n_s^{3D}(0)}{8k_B m \epsilon_c} d$$

A large dielectric constant:
 $\epsilon_c \simeq 90$



Kogan et al., 2009

To understand the large dielectric constant

Kosterlitz's RG equation:

$$\frac{d}{dl}K^{-1}(l) = 4\pi^3 y^2(l),$$

$$\frac{d}{dl}y(l) = [2 - \pi K(l)]y(l)$$

Dielectric constant: $\epsilon_c = \frac{\rho_s^0(T_c^-)}{\rho_s^R(T_c^-)}$


InOx: 1.4-1.9; YBCO: 4.6, 6;
“critical value”: 1.74

Large dielectric constant means large fugacity, or
small vortex core energy.

$$E_c/k_B T_{\text{BKT}} \simeq (A^{1/\theta}/2\pi)\epsilon_c^{-(1-\theta)/\theta}$$

$$\theta = 0.83$$

$$A \simeq 8.62$$




An unsolved problem:


Phase diagram of 2d vortex system at high density

$$\frac{\mathcal{H}_v}{k_B T} = -\pi K \sum_{ij} n_i n_j \log \frac{|\mathbf{r}_i - \mathbf{r}_j|}{R_0} - \log y \sum_i n_i^2$$

Some “contradicting” numerical results:




Lee, Teitel, 1992



Lidmar, Wallin, 1997

Effect of magnetic vortex core



CeCoIn5: close to a magnetic QCP

SC vortices coupled to spin fluctuations:

$$Z[n(\mathbf{r})] = \int \mathcal{D}\Phi^* \int \mathcal{D}\Phi \exp \left(-\frac{\mathcal{H}_v}{k_B T} - \mathcal{S}_{sf} \right)$$

$$\frac{\mathcal{H}_v}{k_B T} = -\pi K \sum_{ij} n_i n_j \log \frac{|\mathbf{r}_i - \mathbf{r}_j|}{R_0} - \log y \sum_i n_i^2$$

$$\mathcal{S}_{sf} = \int d^2\mathbf{r} \int_0^\beta d\tau \left[\frac{1}{2} (\partial_\tau \phi + ig\mu_B \mathbf{H} \times \phi)^2 + \frac{1}{2} (\nabla \phi)^2 + \frac{\alpha}{2} \phi^2 + \frac{\lambda}{4} \phi^4 \right]$$

$$H(\mathbf{r}) = \sum_i n_i H_0(\mathbf{r} - \mathbf{R}_i)$$


Vortex core energy

Free energy density: $\mathcal{F}_\Phi = |\nabla\Phi|^2 + (\alpha - g^2\mu_B^2 H^2(r))|\Phi|^2 + \gamma|\Phi|^4$

$$H_0(\mathbf{r}) \sim (\Phi_0/\lambda^2)K_0(r/\lambda)$$

Spin fluctuations reduce vortex core energy.


$$\delta E_c = \int \tilde{d}^2\mathbf{r} \mathcal{F}[\Phi(\mathbf{r})] \sim -g^4\mu_B^4\tilde{\Phi}_0^4/\gamma\lambda^6 \equiv -V_0 < 0.$$





Magnetic order parameter

Superconducting order parameter

Superfluid density in quantum critical superconductors



Kogan *et al.*, 2009



Hashimoto *et al.*, 2012

Open questions for theorists:

- 1, Phase diagram of vortex system at high density
- 2, BKT transition with competing orders
- 3, Superfluid density in quantum critical superconductors

Is there a holographic solution?